

# Amino acid profile of soybean (Glicine max) sprout protein for determining insulin stimulation amino acids

ORIGINALITY REPORT

# 23%

SIMILARITY INDEX

PRIMARY SOURCES

- |   |   |                |
|---|---|----------------|
| 1 | <a href="https://repository.ubaya.ac.id">repository.ubaya.ac.id</a><br>Internet   | 119 words — 4% |
| 2 | B Kanetro, D H Swasono, T D Astuti, A Sahrah. "Profile of cholesterol and glucose on the various age of volunteers with composite flour diet", IOP Conference Series: Earth and Environmental Science, 2019<br>Crossref | 84 words — 3%  |
| 3 | <a href="https://chemistry.tutorvista.com">chemistry.tutorvista.com</a><br>Internet   | 50 words — 1%  |
| 4 | <a href="https://media.neliti.com">media.neliti.com</a><br>Internet   | 42 words — 1%  |
| 5 | J. Derek Bewley, Michael Black. "Chapter 6 Mobilization of Reserves", Springer Science and Business Media LLC, 1978<br>Crossref   | 41 words — 1%  |
| 6 | Bayu Kanetro. "Hypocholesterolemic Properties of Protein Isolate from Cowpeas (Vigna Unguiculata) Sprout in Normal and Diabetic Rats", Procedia Food Science, 2015<br>Crossref  | 37 words — 1%  |
| 7 | <a href="https://ifrj.upm.edu.my">ifrj.upm.edu.my</a><br>Internet   | 29 words — 1%  |
| 8 | <a href="https://repository.wima.ac.id">repository.wima.ac.id</a><br>Internet   | 28 words — 1%  |