

Prosiding IFC 2016_UNIV WIDYA MANDALA SURABAYA-304-310.pdf

The Physical Properties and Acceptability of Duck Meat Cured in Curcumin Extract with Added STPP

Sri Hartati Candra Dewi*, Chatarina Wariyah, and Niken Astuti

¹ Department of Husbandry, Faculty of Agroindustry, Mercuru Buana Yogyakarta University, Yogyakarta, 55753, Indonesia

Abstract

Duck meat was taken from laying ducks which had become unproductive and also from older male ducks. The duck meat texture was tough and the high fat content of the duck meat caused it to oxidize easily during storage and processing. It is known that curing duck meat in curcumin extract and freezing for storage can inhibit oxidation of the fat, but the meat turns yellowish in colour and the texture of the frozen meat when cooked is tougher. The purpose of this study was to evaluate the effect of curing of duck meat in curcumin extract with the addition of STPP (sodium tripolyphosphate) on the physical properties (colour, texture and water holding capacity) of the duck meat and on its acceptability. The research used Randomized Complete Design, and the duck meat was cured in curcumin extract at concentrations of 0.3% and 0.4% (w/w), then STPP was added at various concentrations of: 0.0; 0.10; 0.15; 0.20 and 0.25% (w/w). The duck meat texture was tested using a Texture Analyzer and the colour tested with a Colour Reader. Acceptability of the cured duck meat with added STPP was determined by Hedonic Test based on smell, colour, texture and flavour. The results of this study concluded that raw duck meat texture was based on resistance to force and deformation, which were not significantly different. Resistance to force was between 545,91g and 597,76g and the deformation was between 70,47% and 73,03%. Water holding capacity of cured meat was between 40,05% and 46,77%, which is higher than fresh meat at 26,81%. However, the color of the cured duck meat wasn't as bright as fresh duck meat, as shown by the lightness (L) of between 34,04 and 36,46 against 34,43 for the fresh duck meat, and yellowness (b) of between 21,43 and 22,98 for cured meat and 22,49 for fresh meat. The research showed that the duck meat cured with curcumin extract at 0.3% and 0.4% concentrations and added with 0:10 to 0:20% STPP resulted in the most acceptable product, based on texture, color and water holding capacity, especially the brighter color and softer texture.

Keywords: duck meat, curcumin, STPP, texture.

Introduction

Duck (*Anas platyrhynchos*) is a type of poultry farmed for meat and eggs. The quantities of duck meat on the market are still very limited as supply mostly comes from culled females (54.35%), but as much as 35.41% can come from male salvage and up to 18% from young females (Hardjosworo, 2001). Duck meat is the meat of culled non-productive female layers and older males. Reject duck meat has a clay-like texture and a fat content reaching 1.84%, in contrast to chicken meat at only 1.05% fat (Ali *et al.*, 2007). Unsaturated fatty acids (AL TJ) make up more than 60% of the total fatty acids, which results in the duck meat being easily oxidized, thereby degrading flavour, destroying nutrients and leading to a build-up of toxic substances. According to Baggio and Bragagnolo (2006) meat, during processing and storage, can be the subject of oxidation induced by the presence of heat, light, metal and oxygen which will produce ROS (Reactive Oxygen Species) such as aldehydes, peroxide and cholesterol oxides that can lead to degenerative diseases such as cardiovascular disease and early aging.

Attempts to inhibit oxidation of the fat in duck meat were made by Dewi and Astuti (2013) by adding 0.3% turmeric extract as a natural source of antioxidants and then curing for 10 minutes. Storage was undertaken for 8 weeks in a freezer. Turmeric is known to contain curcumin which can inhibit lipid peroxidation (Jayaprakasha *et al.*, 2006). The results show that extract of turmeric can inhibit an increase in the numbers of peroxides and TBARS of duck meat, and the texture of duck meat after storage becomes more tender. But curcumin is yellow in colour and has a distinctive turmeric flavour,

which can affect the acceptability of the product. In addition, the research is still limited to storage only. Yet according to Sampaio *et al.* (2012), lipid oxidation will continue during cooking.

The problem is that the texture of duck meat becomes more tender after storage in a freezer (Dewi and Astuti, 2013), but according to Fernandes *et al.* (2013) storing lamb and mutton at freezing temperatures (-18° C) causes low water retention or Water Holding Capacity (WHC), so that the texture of the meat after cooking is hard. Abdel *et al.* (-) states that the addition of sodium tripolyphosphate (STPP) to lamb meat which is to be frozen can inhibit the decrease in WHC compared to the control, so the texture of the cooked meat is softer and is preferred by the consumer. And according to Marsha *et al.* (2013) the use of STPP in turkey meat, in addition to controlling the WHC, can also inhibit the oxidation of fat by slowing down the penetration of heat into the material. Hence this objective study to evaluate the effects of curing duck meat in curcumin extract, and the addition of STPP on the physical properties and acceptability of duck meat, both raw and after cooking.

Materials and Methods

This study was undertaken with the aim of producing duck meat with a soft texture after frozen storage, to measure the effects of antioxidants, provide high acceptability and stability during processing, and to be safe for human consumption.

Materials

The materials used for the study of duck meat were derived from duck breeders in the village of Argomulyo, Sedayu, Bantul, Yogyakarta. Turmeric, as a natural source of antioxidants, was purchased from a local market in the Yogyakarta area. Analysis of the base material (duck meat) included water content (AOAC, 1990). Water holding capacity (Hamm, 1960 in Soeparno, 2009) was tested using a Texture Analyzer and colour was tested using a Colour Reader. Chemicals used in all the pro qualifying analyses were obtained from Merck.

Methods

The research method consists of five steps. These are : 1) Preparation of turmeric curcumin extract by sorting tubers, then peeling and washing. Curcumin extraction using maceration method (Marsono *et al.*, 2005), 2) Curing fresh duck meat with turmeric curcumin extract (with a variation of 0.3% and 0.4%) and variation of the addition of STPP at 0.00; 0.10; 0.15; 0.20; and 0.25%, 3) Storage of duck meat (phase 2) at freezing temperatures (-10°C) for 8 weeks, 4) Testing the physical properties (texture, color, WHC), 5) Determining the organoleptic acceptability of (phase 3) raw duck meat by sensory test.

3 Statistical analysis

The experimental design used was completely randomized design, factorial pattern with factors such as variation of the ratio of fresh meat to the amount of curcumin extract and STPP to determine the differences between treatments used by the F test, then the real difference between the samples was determined by Duncan's Multiple Range Test (DMRT) (Gacula and Singh, 1984)

Result and discussion

Texture of duck meat

The results showed that there was no interaction effect from each curcumin and STPP treatment on the texture of the cured duck meat (Table 1). The addition of 0.3% and 0.4% curcumin did not affect the texture of the cured raw duck meat. This is consistent with the results of research by Dewi and Astuti (2014) which states that adding curcumin only increases antioxidants, which does not affect the texture of duck meat. The texture of the cured raw duck meat is also not influenced by the addition of STPP because STPP, as well as increasing the water retention capabilities of meat during cooking, can also help maintain meat texture and tenderness (Petracci *et al.* 2013). Tender meat has a much higher market value than tough meat. This is likely due to the meat used being a sample of different muscles, as in this study by using only a few ducks but utilizing all the meat from each carcass. Soeparno (2009) states that muscle difference affects the texture and tenderness of meat.

Table 1. Texture of cured duck meat (g)

Curcumin (%)	STPP (%)					Average (ns)
	0	0,1	0,15	0,2	0,25	
0,3	559,25	542,50	686,50	599,88	578,38	593,30
0,4	565,75	650,50	509,01	491,94	554,88	554,42
Average (ns)	562,50	596,50	597,76	545,91	566,63	

Deformation of duck meat

Deformation is a change in the meat texture caused by cutting the meat to measure texture. Raw cured duck meat has relatively similar deformation and is not affected by curcumin treatment, STPP and interaction (Table 2). This may be because the texture is not significant so deformation is insignificant as well.

Table 2. Deformation of cured duck meat (%)

Curcumin (%)	STPP (%)					Average (ns)
	0	0,1	0,15	0,2	0,25	
0,3	55,44	77,63	64,51	72,65	72,22	68,49
0,4	69,57	68,43	76,42	69,32	72,00	71,15
Average (ns)	62,51	73,03	70,47	70,99	72,11	

If the cured duck meat was overcooked, the deformation was affected by the addition of STPP. Deformation of cured cooked duck meat was greatest with the addition of STPP by 0.1%. Higher additions of STPP showed no more significant deformation. This is because in meat products, STPP is generally used to maintain texture or tenderness after cooking.

Brightness (L) of cured duck meat

The brightness of cured duck meat was influenced significantly by the addition of STPP, but there was no significant effect from the addition of curcumin as well as its interaction (Table 3). The addition of 0.4%

curcumin gave the same relative brightness as the addition of 0.3%. Curcumin only serves to add antioxidants, which are substances that can inhibit an oxidation reaction in materials susceptible to oxidation (Fennema, 1996). It is likely that it has no real effect on the brightness of the flesh. The brightness of cured duck meat was significantly enhanced by the STPP treatment with the lowest brightness coming from the addition of 0.2% STPP and the highest brightness of 36.46 from 0.1% addition of STPP. Petracci *et al.* (2013), stated that some of the functional ingredients used in meat, STPP being one of them, are also able to increase the water retention of meat during cooking.

Table 3. Lightness of cured duck meat

Curcumin (%)	STPP (%)					Average(ns)
	0	0.1	0.15	0.2	0.25	
0.3	35,08	36,17	35,82	30,35	33,73	34,20
0.4	33,78	36,74	36,45	32,99	34,34	34,85
Average	34,43 ^a	36,46 ^a	36,14 ^a	31,67 ^b	34,04 ^a	

Redness of cured duck meat

The colour of cured duck meat is not affected by the addition of curcumin, STPP or its interaction (Table 4). Redness value which is positive showed that duck has a reddish colour, while if the value is negative then the meat will be a greenish colour. In this study the positive value means that the cured duck meat in this research is reddish in colour. Red cured duck meat indicates normal meat.

Table 4. Redness of cured duck meat

Curcumin (%)	STPP (%)					Average (ns)
	0	0,1	0,15	0,2	0,25	
0.3	7,58	6,25	7,00	7,67	6,22	6,94
0.4	7,20	8,07	7,38	6,69	7,02	7,27
Average (ns)	7,39	7,16	7,19	7,18	7,52	

Yellowish color of cured duck meat

The result showed that the value of yellowness of cured duck meat was significantly influenced by the addition of curcumin, but was not influenced by the addition of STPP and there was no interaction (Table 5). In addition, curcumin added at 0.4% had a higher yellowness value, i.e. had a more yellow coloured flesh than the 0.3%.

Table 5. Yellowness of cured duck meat

Curcumin (%)	STPP (%)					Average
	0	0,1	0,15	0,2	02525	
0.3	20,17	21,40	24,05	21,26	21,76	21,73 ^a
0.4	24,80	24,10	20,40	21,60	24,20	23,02 ^b
Average (ns)	22,49	22,75	22,23	21,43	22,98	

5

Ali *et al.* (2007) stated that the color of the duck meat has a very high redness value, but it has a low brightness value. Dewi and Astuti (2014) stated that the addition of curcumin gave an improved flesh color. It is further mentioned that there is a relationship between the amount of curcumin and the length of time curing the duck meat on the colour. i.e. if lower levels of curcumin are used to get the desired yellow color, then curing time has to be longer.

Water Holding Capacity (WHC) Value of cured duck meat

The results of this study indicate that the WHC value of duck meat was significantly influenced by the addition of STPP, but that it was not affected by the addition of curcumin and its interaction (Table 6).

The addition of STPP increased the WHC value, but at an addition of 0.25% (Table 6).

Table 6. WHC Value of cured duck meat

Curcumin (%)	STPP (%)					Average (ns)
	0	0,1	0,15	0,2	0,25	
0,3	30,28	41,01	43,00	43,22	38,82	39,27
0,4	23,33	41,25	50,13	50,32	41,28	41,26
Average	26,81 ^a	41,13 ^b	46,57 ^b	46,77 ^b	40,05 ^b	

This is consistent with the statement of Thomas (1997) cit. Dewanti (2009) that STPP can absorb, bind and hold water, increase water holding capacity (WHC) and tenderness. Yuanita (2008) states that the STPP FG is able to break the bonds of the water in the meat, thus causing a decrease in the levels of WA (water activity) which is an essential component in inhibiting the growth of microbes. A high WHC value for meat shows that the meat can hold water, so the water level is higher. This is consistent with the research results of Pratiwi (2016), which state that higher the levels of STPP FG used, up to a level of 4% in broiler meat soaking, the lower the water content. An improvement in the FG STPP treatment levels is also accompanied by an increase in the elasticity of the meat and its water holding capacity. This means that a decrease in water content does not occur in the water-bound or immobilized free water, but is due to the evaporation of free water (Yuanita *et al.* 1997).

Sensory Test of Cured Duck Meat

The results of the study of sensory testing of cured raw duck meat show that the smell, texture are affected on the whole by the addition of curcumin and STPP (Table 7). The smell of the raw meat with curcumin added at 0.3% is less preferred than that at 0.4%;, with the highest score of unlikeability being 0.3% curcumin and 0.2% STPP giving a score of 4.60 or the most disliked. The colour test by sensory testing was not affected by either the curcumin treatment or the STPP. The texture of the raw duck meat was tested by pressing with a finger. The results of the tests of sensory texture of the meat turned out to be significantly different, with the least preferred texture coming from the addition of 0.4% curcumin and 0.25% STPP. But the most preferred overall by the panelists in a sensory test came from the addition of curcumin at 0.3% and 0.4% and STPP 0.25%.

In the sensory test of cured cooked duck meat the smell was not affected by treatment with curcumin and STPP. While colour, texture and taste were on the whole affected by treatment with curcumin and STPP,

the addition of curcumin made no significant difference to the smell of the cooked meat, possibly because 0.3% and 0.4% additions of curcumin after cooking had a relatively similar smell.

Table 7. Sensory Test of Cured Duck Meat

		Smell	Color (ns)	Texture	Overall
Basic Material		3,53 ^{ab}	3,47	4,27 ^b	3,73 ^{ab}
Curcumin (%)	STPP (%)				
0,30	0,00	4,53 ^{ab}	3,73	4,20 ^{ab}	4,33 ^b
	0,10	4,27 ^{ab}	3,21	3,33 ^{ab}	3,73 ^b
	0,15	3,80 ^{ab}	3,33	3,27 ^{ab}	3,47 ^{ab}
	0,20	4,60 ^b	4,00	3,87 ^{ab}	4,13 ^{ab}
	0,25	4,47 ^{ab}	3,40	3,33 ^{ab}	3,25 ^{ab}
0,4	0,00	3,30 ^a	3,67	3,47 ^{ab}	3,53 ^{ab}
	0,10	3,73 ^{ab}	3,40	3,93 ^{ab}	3,40 ^a
	0,15	3,33 ^a	3,47	3,93 ^{ab}	3,47 ^{ab}
	0,20	4,13 ^{ab}	3,60	3,67 ^{ab}	3,87 ^{ab}
	0,25	3,40 ^{ab}	3,53	3,07 ^a	3,20 ^a

Table 8. Sensory Test of Cured Duck Meat

		Smell (ns)	Color	Texture	Flavore	Whole
Basic Material		3,67	4,40 ^b	3,13 ^{ab}	3,33 ^{ab}	3,53 ^{ab}
Curcumin (%)	STPP (%)					
0,30	0,00	3,20	3,80 ^{ab}	3,80 ^{ab}	4,33 ^{bcd^{ef}}	4,20 ^{abc}
	0,10	3,33	3,33 ^{ab}	3,19 ^{ab}	3,13 ^a	3,47 ^a
	0,15	3,93	3,60 ^{ab}	2,73 ^a	3,93 ^{abc^{de}}	3,73 ^{ab}
	0,20	3,27	3,13 ^a	2,80 ^a	3,47 ^{abc}	3,30 ^a
	0,25	3,67	3,44 ^{ab}	3,47 ^{ab}	3,80 ^{abcd}	3,93 ^{abc}
0,4	0,00	3,13	3,27 ^a	4,20 ^{ab}	4,67 ^{def}	4,13 ^{abc}
	0,10	4,20	3,13 ^a	4,27 ^b	5,00 ^{ef}	5,00 ^c
	0,15	4,04	3,47 ^{ab}	4,33 ^b	5,13 ^f	4,67 ^{bc}
	0,20	4,07	3,07 ^a	3,53 ^{ab}	4,60 ^{c^{def}}	4,40 ^{abc}
	0,25	4,13	3,40 ^{ab}	3,20 ^{ab}	4,60 ^{c^{def}}	4,40 ^{abc}

While the most preferred colour of cooked cured meat was achieved by adding STPP at 0.2% and curcumin at 0.3% to 0.4%. Preferred texture of the cooked cured meat is achieved by addition of curcumin 0.3% and STPP at 0.1 to 0.2%, and curcumin at 0.4% with STPP at 0.2% to 0.25%. While the preferred taste of cooked cured meat is with curcumin added at 0.3% and STPP from 0.1% to 0.25%, and curcumin at 0.4% with STPP from 0.2% to 0.25%. On the whole the most preferred cooked cured duck meat is with an addition of curcumin at 0.3% with STPP from 0.1% to 0.2%, and curcumin added at 0.3% with STPP at 0.20% to 0.25%.

Conclusion

The results of this study concluded that raw duck meat texture was based on resistance to force and deformation, which were not significantly different. Resistance to force was between 545,91g and 597,76g and the deformation was between 70,47% and 73,03%. Water holding capacity of cured meat was between 40,05% and 46,77%, which is higher than fresh meat at 26,81%. However, the color of the

cured duck meat was not as bright as fresh duck meat, as shown by the lightness (L) of between 34,04 and 36,46 against 34,43 for the fresh duck meat, and yellowness (b) of between 21,43 and 22,98 for cured meat and 22,49 for fresh meat. The research showed that the duck meat cured with curcumin extract at 0.3% and 0.4% concentrations and added with 0:10 to 0:20% STPP resulted in the most acceptable product, based on texture, color and water holding capacity, especially the brighter color and softer texture.

1 **knowledge**

We gratefully acknowledge the Directorate of Research and Community Service, Directorate General of Reinforcement and Development, Ministry of Research, Technology and Higher Education of the Republic of Indonesia for providing financial support from the Competitive Research Grant Program in 2016.

References

- Abdel-Ad Aal, H.A. and H.M.A. Mohamed. -. Change in the ultrastructure of lamb muscles as influenced by freezing and sodium tripolyphosphate (STPP). Department of Food Science, Faculty of Agriculture, Minia University, El-Minia, Egypt. *Lucransttiitific*. Vol. 56. SeriaZootehnie.
- Ali. M.S., G.H. Kang, H.S. Yang, J.Y. Jeong, Y.H. Hwang, G.B. Park dan S.T. Joo. 2007. A Comparison of meat characteristics between duck and chicken breast. *Asian-Aust. J. Anim. Sci.* 20 : 1002-1006.
- AOAC, 1990. *Official Methods of Analysis Association Official Agricultural Chemistry*. Washington D.C.
- Baggio, S.R. dan N. Braganolo, 2006. Cholesterol oxide, cholesterol, total lipid and fatty acid content in processed meat products during storage. *J. Food Sci.* 71 : 513-520.
- Dewanti, T. 2009. STPP Pengganti Boraks (Bleng) pada Krupuk Puli dan Bakso, <http://terminalcurhat.blogspot.com/2009/03/stpppengganti-boraks-bleng-pada-krupuk.html> diakses 6 Januari 2016
- Dewi, S.H.C. 2010. Sifat Fisik Daging Sapi pada Lama Penyimpanan yang Berbeda. Laporan Penelitian. Universitas Mercu Buana. Yogyakarta.
- Dewi, S.H.C. 2011. Populasi Mikroba dan Sifat Fisik Daging Sapi Beku pada Lama Penyimpanan yang Berbeda. Laporan Penelitian. Universitas Mercu Buana. Yogyakarta.
- Dewi, S.H.C. dan Astuti, N. 2013. Ekstrak Kurkumin Kuyut Untuk Menghambat *Off-Flavor* Daging Itik Selama Pangolahan. Laporan Penelitian. LPPM. Universitas Mercu Buana Yogyakarta.
- Dewi, S.H.C. dan Astuti, N. 2014. Akseptabilitas dan Sifat Daging Itik Afkir yang Dilakukan Curing Menggunakan Ekstrak Kurkumin Kuyut untuk Menghambat Oksidasi lemak Selama Penyimpanan. *Jurnal Agritech*. Fakultas Teknologi Pertanian. Universitas Gadjah Mada. Yogyakarta. Vo. 34 (4) : 415-421.
- Fennema, O.R., 1996. *Principles of Food Science*. Marcell Dekker Inc. New York.
- Fujiwara, H., M. Hosokawa, X. Zhou, S. Fujimoto, K. Fukuda, K. Toyoda, Y. Nishi, Y. Fujito, K. Yamada, Y. Yamada., Y. Seino and N. Inagaki. 2008. Curcumin Inhibits Glucose Production in Isolated Mice Hepatocytes. *Diabetes Research and Clinical Practice*. 80 : 188-191.
- Gacula, M.C. dan J. Singh, 1984. *Statistical Methods in Food and Consumer Research*. Academic Press, Inc. Orlando. San Diego. New York. London.
- Hardjosworo. 2001. *Ternak Itik*. Penebar Swadaya. Jakarta.
- Jayaprakasha, G.K, L. Jaganmohan Rao , K.K. Sakariah. 2006. Antioxidant activities of curcumin, demethoxy curcumin and bisdemethoxy curcumin *Food Chemistry* 98 : 720-724.
- Marsha, L.S., J.R. Clause, N.G. Marriots, W.N. Eigel, H. Wang. 2013. Reduction in Lipid Oxidation by Incorporation of Encapsulated Sodium Tripolyphosphate in Ground Turkey. *Meat Science*. 95: 376-380.
- Marsono, Y., Safitri, R. dan Noor, Z., 2005. Antioksidan dalam Kacang-kacangan : Aktivitas dan Potensi serta Kemampuannya Menginduksi Pertahanan Antioksidan pada Model Hewan percobaan. Laporan Penelitian Hibah Bersaing XII.
- Petracci, M., M. Bianchi, S. Mudalal, C. Cavani. 2013. Functional Ingredients for Poultry Meat Products. *Trend in Food Science and Technology*. 33 : 27-39.
- Pratiwi, Y.D. 2016. Pengaruh kadar *sodium tripolyphosphate food grade* (STPP FG) terhadap kualitas daging broiler. Skripsi. Universitas Mercu Buana Yogyakarta. Yogyakarta.
- Sampaio, G.r., T. Saldanna, R.a.M. Soares, E.A.F.S. Torres. 2012. Effect of natural Antioxidant Combinations on Lipid oxidation in Cooked Chicken Meat During Refrigerated Storage. *Food Chemistry*. 135 : 1383-1390.
- Soepamo. 2009. *Ilmu dan Teknologi Daging*. 4th edition. GadjahMada University Press. Yogyakarta.
- Yuanita L., Surodjo S., dan Wikandari P. 1997. *Pengaruh Penggunaan Alkali Fosfat Sebagai Pengganti Boraks Terhadap Kualitas Daging Olahan*. Laporan Penelitian. Lembaga Penelitian IKIP Surabaya.
- Yuanita, L. 2008. Penentuan Kadar STPP *Food Grade* Untuk Meningkatkan Masa Simpan Ikan Nila Tilapia (*Oreochromis niloticus* L.). *Jurnal Berkala Penelitian Hayati*: 13 (179-186).

5%

SIMILARITY INDEX

PRIMARY SOURCES

- | | | |
|---|---|-----------------|
| 1 | www.myfoodresearch.com
Internet | 67 words — 2% |
| 2 | repository.unpas.ac.id
Internet | 17 words — < 1% |
| 3 | baadalsg.inflibnet.ac.in
Internet | 13 words — < 1% |
| 4 | zombiedoc.com
Internet | 10 words — < 1% |
| 5 | J.H. Choi, Y.S. Choi, H.W. Kim, D.H. Song, C.J. Kim.
"Effects of postmortem temperature on the physicochemical characteristics of prerigor Pekin duck breast muscles", Poultry Science, 2016
Crossref | 9 words — < 1% |
| 6 | Figueirêdo, Bruno Chacon, Isabela Jorge Trad, Lilian Regina Barros Mariutti, and Neura Bragagnolo.
"Effect of annatto powder and sodium erythorbate on lipid oxidation in pork loin during frozen storage", Food Research International, 2014.
Crossref | 9 words — < 1% |
| 7 | id.scribd.com
Internet | 8 words — < 1% |
| 8 | K.E. Hwang, H.W. Kim, Y.S. Choi, S.Y. Lee, E.J. Yeo, Y.K. Ham, S.M. Choi, M.A. Lee, C.J. Kim.
"Evaluation of the antioxidant effect of ganghwayakssuk (Artemisia | 8 words — < 1% |

princeps Pamp.) extract alone and in combination with ascorbic acid in raw chicken patties", Poultry Science, 2013

Crossref

9

sinta.ristekbrin.go.id

Internet

8 words — < 1%

10

Kurnia Ramadhan, Nurul Huda, Ruzita Ahmad. "Duck Meat Utilization and the Application of Surimi-like Material in Further Processed Meat Products", Journal of Biological Sciences, 2010

Crossref

7 words — < 1%

11

Ke Li, Lin Chen, Ying-Ying Zhao, Yu-Pin Li, Na Wu, Hao Sun, Xing-Lian Xu, Guang-Hong Zhou. "A comparative study of chemical composition, color, and thermal gelling properties of normal and PSE-like chicken breast meat", CyTA - Journal of Food, 2014

Crossref

6 words — < 1%

EXCLUDE QUOTES OFF

EXCLUDE BIBLIOGRAPHY ON

EXCLUDE MATCHES OFF